

Charu Tiwari, Hemanshi Shah, Jayesh Desale, Mukta Waghmare

NEONATAL GARTNER DUCT CYST: TWO CASE REPORTS AND LITERATURE REVIEW

Dept of Paediatric Surgery, TNMC & BYL Nair Hospital, Mumbai Central,
Mumbai, Maharashtra. India. Pin: 400008

Abstract

Vaginal cysts are rare, particularly in the newborn. They usually present as one of these three entities in the newborn: paraurethral cysts (Skene duct cysts), Gartner duct cysts (mesonephric ductal remnants) or a covered ectopic ureter. Abdominal ultrasound should always be included in the clinical evaluation in search of renal anomalies. We report two cases of Gartner cysts in neonates.

Key words: Neonate, Vaginal Mass, Gartner duct cyst

DEV PERIOD MED. 2017;XXI,1:35-37

INTRODUCTION

Vaginal wall cysts are rare in common urological practice, especially in neonates [1]. Gartner duct cysts (GDC) arise as a consequence of obstruction of the Gartner duct (mesonephric remnant). They are located in the anterior or lateral wall of the vagina [1, 2]. They may be associated with renal and ureteral anomalies [1]. The correct treatment is the removal of the entire cyst through a vaginal approach [1]. Aspiration, marsupialisation and deroofting are the various treatment options [1]. Accurate diagnosis can only be made by histology. We report two neonates with Gartner duct cysts aiming to emphasize their embryogenesis, clinical presentations along with differential diagnoses, work-up and management.

CASE STUDIES

Case 1: A 5-day-old healthy female neonate was admitted with complaints of an asymptomatic vaginal mass, noticed by her mother. General clinical examination was essentially within normal limits. Perineal examination showed normal urethral, vaginal and anal openings. A cystic lesion arising from the right lateral vaginal wall could be seen (Fig. 1).

Abdominal ultrasound was normal. On cystogenitoscopy, the urethra, bladder and both ureteral openings were normal. Vaginoscopy confirmed the cystic lesion in the right lateral vaginal wall. The cyst was deroofted. Histopathology revealed Gartner duct cyst. The patient is asymptomatic on follow-up.

Case 2: A 6 day-old female neonate was admitted with a swelling over her back. On examination, she had a large lumbosacral rachischisis along with paraplegia and incontinence of the bladder and bowel. On clinical examination of the perineum, the introitus was normal with urethral and vaginal openings. A cystic mass was noted in the left anterolateral wall of the vagina (Fig. 2). The anus was patulous.

Abdominal ultrasound revealed an absent left kidney. On needle aspiration, white milky fluid was aspirated and the cyst collapsed, suggesting Gartner duct cyst.

DISCUSSION

GDCs represent cystic remnants of the wolffian (mesonephric) duct system and contribute approximately 11% of all vaginal cysts [3]. These develop as a result of incomplete regression of the mesonephric or wolffian duct during fetal development [4]. They are located submucosally along the anteromedial wall of the vagina [2, 4]. Although usually solitary, they may be multiple [4, 5]. The cysts are usually small, less than 2cm in diameter; however, giant cysts have also been reported [5]. The cyst wall is lined with a non-mucin-secreting, squat, cuboidal epithelium [3]. The fluid within the cyst is white, thick, and viscous [3]. Mucoïd fluid rules out a mesonephric cyst [3].

If there is a history of antenatal exposure to synthetic hormones, adenosis of the vagina must be considered [4]. The presence of mucosa, which normally stains with Lugol's solution, helps to exclude the diagnosis of adenosis [4].



Fig. 1. A view of the cystic lesion arising from the right lateral vaginal wall



Fig. 2. A view of the cystic mass in the left anterolateral wall of the vagina.

GDC is extremely rare in the paediatric age group and more so in neonates [6]. The differential diagnosis of Gartner cyst in a neonate is broad [2]. It includes imperforate hymen (most common), paraurethral Skene's duct cyst, urethral prolapse, prolapsed ectopic ureterocele, urethral polyp, congenital lipoma, vaginal

prolapse, rhabdomyosarcoma of the vagina [1, 7]. Vaginal inclusion cysts and Bartholin gland cysts are usually seen in the third and fourth decades and not in neonates [4].

Accurate diagnosis requires a thorough understanding of the diagnostic possibilities and a systemic evaluation [2]. Physical examination remains the most useful tool for determining the specific pathology [2]. During examination, the patient should be placed in frog-leg position and the labia majora should be gently grasped and pulled caudally and laterally enabling funneling of the introitus and vagina [1]. The size of the clitoris, hymenal configuration, urethral location and the site and character of the mass should be carefully noted [2]. Placing a small feeding tube within the suspected urethral orifice can further help in local examination [2]. Abdominal ultrasound (USG) is a helpful adjunct in diagnosing associated renal disorders [2].

The histology provides the accurate diagnosis on the basis of embryological origin [7]. The vagina is derived from the paramesonephric (Mullerian) duct, the mesonephric (Wolfian) ducts and the urogenital sinus [7]. The vaginal cysts are lined with stratified squamous epithelium as they originate from the Mullerian duct. Gartner duct cysts are lined with cuboidal epithelium {mesonephric/Wolfian origin} [7]. Transitional epithelium in the cyst wall confirms the origin to be in the urinary tract (Skene's ducts cysts, paraurethral cysts, urethral prolapsed, ureterocele prolapsed and ectopic ureter) and calls for complete renal work-up [7, 8, 9].

Association of GDC with ipsilateral renal agenesis or dysplasia is rare and is caused by the abnormal development of the ureter [10, 11]. The presence of ureteric ectopia associated with GDC has been reported to be caused by the failure of separation of the ureteric bud from the mesonephric duct, which leads to persistence of Gartner's duct, frequently with cystic dilation [10, 11]. Presentation with sepsis or non- or poorly functioning renal tissue, is an indication for ureterectomy or nephroureterectomy on the affected side [10].

Management options are aspiration, deroofing, marsupialisation and complete excision [1, 4, 5, 9, 12]. A recent study by Rios et al involving four women has shown that conservative treatment can be a safe option for asymptomatic patients with vaginal GDCs [13]. In a study involving 15 patients by Abd-Rabbo et al, aspiration and injection sclerotherapy with Tetracycline has been reported as one of the management options [14]. This technique has been reported as an ideal, safe and effective simple office procedure for management of symptomatic Gartner cysts [abd14]. However, its application in neonates is unknown. Reassuringly, the long-term prognosis is excellent [12].

CONCLUSION

Neonatal Gartner Duct cysts are rare. Association with renal anomalies needs to be ruled out during evaluation. Management is simple and long-term prognosis is excellent.

REFERENCES

1. Molina Escudero R, Navas Martinez M-C, Castillo O-A. Vaginal Gartner cysts: clinical report of four cases and a bibliographic review. *Arch Esp Urol*. 2014;67(2):181-184.
2. Rink R-C, Kaefer M. Surgical management of disorders of sexual differentiation, cloacal malformation, and other abnormalities of external genitalia in girls. In: Wein AJ editor. *Campbell-Walsh Urology*. 9th ed. Philadelphia: Saunders; 2007;3629-3666.
3. Bats A-S, Metzger U, Le Frere-Belda M-A, Brisa M, Lecuru F. Malignant transformation of Gartner cyst. *Int J Gynecol Cancer*, 2009;19:1655-1657.
4. Delmore J, Glob. Benign neoplasms of the vagina. *Libr. Women's Med.* (ISSN: 1756-2228) 2008.
5. Vlahovic A, Stankovic Z-B, Djuricic S, Savic D. Giant Gartner duct cyst and elevated CA-125. *J Pediatr Adolesc Gynecol*. 2014;27(6):e137-138.
6. Paradies G, Zullino F, Caroppo F, Orofino A, Lanzillo M-P, Leggio S. Gartner's duct cyst: report of three cases. (in Italian) *Pediatr Med Chir*. 2011;33:247-252.
7. Moralioglu S, Bosnalı O, Celayir A-C, Şahin C. Paraurethral Skene's duct cyst in a newborn. *Urol Ann*. 2013;5(3):204-205.
8. Bergner DM. Paraurethral cysts in the newborn. *South Med J*. 1985;78:749-750.
9. Kimbrough HM Jr, Vaughan ED Jr. Skene's duct cyst in a newborn: case report and review of the literature. *J Urol*. 1977;117:387-388.
10. Holmes M, Upadhyay V, Pease P. Gartner's duct cyst with unilateral renal dysplasia presenting as an introital mass in a new born. *Pediatr Surg Int*. 1999;15:277-279.
11. Dwyer PL, Rosamilia A. Congenital urogenital anomalies that are associated with the persistence of Gartner's duct: a review. *Am J Obstet Gynecol*. 2006;19:354-359.
12. Binsaleh S, Al-Assiri M, Jednak R, El-Sherbiny M. Gartner duct cyst simplified treatment approach. *Int Urol Nephrol*. 2007;39(2):485-487.
13. Rios SS, Pereira LCR, Santos CB, Chen ACR, Chen JR, Vogt MB. Conservative treatment and follow-up of vaginal Gartner's duct cysts: a case series. *Journal of Medical Case Reports* 2016;10:147.
14. Abd-Rabbo MS, Atta MA. Aspiration and tetracycline sclerotherapy: a novel method for management of vaginal and vulval Gartner cyst. *Int J Gynecol Obstet*. 1991;35:235-237.

Acknowledgement: Nil.

Source of Support: Nil

Conflicts of Interest: Nil

This manuscript has been read and approved by all the authors, the requirements for authorship have been met and each author believes that the manuscript represents honest work.

Received/Nadesłano: 17.01.2017 r.

Accepted/Zaaakceptowano: 22.02.2017 r.

Published online/Dostępne online

Address for correspondence:

Hemanshi Shah

Dept of Paediatric Surgery, TNMC & BYL Nair Hospital,
Mumbai Central, Mumbai, Maharashtra.

India. Pin: 400008.

Contact No. 022-23027324, Fax No. 0712-6631896

e-mail ID: hemanshisshah@gmail.com

First Author:

Charu Tiwari

TNMC & BYL Nair Hospital,

Mumbai, Maharashtra

India Pin: 400008

e-mail ID: drcharusharma18@gmail.com