

Joanna Gajewska¹, Jadwiga Ambroszkiewicz¹, Halina Weker², Magdalena Chęłchowska¹

RELATIONS BETWEEN OXIDIZED LOW-DENSITY LIPOPROTEINS AND FAT-SOLUBLE VITAMIN CONCENTRATIONS IN OBESE CHILDREN – PRELIMINARY STUDY

ZALEŻNOŚCI POMIĘDZY STĘŻENIEM OKSYDOWANYCH LIPOPROTEIN NISKIEJ GĘSTOŚCI A STĘŻENIEM WITAMIN ROZPUSZCZALNYCH W TŁUSZCZACH U DZIECI OTYŁYCH – DONIESIENIE WSTĘPNE

¹Screening Department

²Department of Nutrition

Institute of Mother and Child in Warsaw, Poland

Abstract

Introduction: Although lipid peroxidation products are formed during normal cell metabolism, they appear mostly in pathological conditions via producing an excess of free radicals that can react with unsaturated fatty acids, in particular low-density lipoprotein (LDL).

The aim of this study was to investigate the relations between oxidized LDL (oxLDL), the fat-soluble vitamin status and the anthropometric parameters in prepubertal obese children.

Material and methods: Thirty-two obese (SDS-BMI >2) and 25 non-obese children (SDS-BMI <-1+1>) were included in the study. The concentration of oxLDL was determined in the serum by the ELISA assay. Vitamin A and E were measured by the high-pressure liquid chromatography method. Total cholesterol, LDL- and HDL-cholesterol, and triglyceride levels were determined by enzymatic methods.

Results: The concentrations of oxLDL and vitamin A were higher in obese children than in normal-weight controls by about 50% ($p=0.01$) and 40% ($p=0.001$), respectively. In obese children the significant positive correlation was found between oxLDL and vitamin A concentrations ($p<0.05$). In addition, oxLDL correlated positively with BMI values ($p<0.05$) and the amount of fat mass (kg) ($p<0.02$) in these patients. Concentrations of vitamins A and E correlated with the level of total cholesterol ($p<0.05$; $p<0.01$, respectively). Moreover, a positive correlation between vitamin E and LDL-cholesterol was observed ($p<0.05$).

Conclusions: Our preliminary study shows that oxLDL starts early during the prepubertal period and may precede atherosclerotic lesions. We suggest there is an occurrence of relationships between vitamin A and oxidized LDL in prepubertal obese children. Vitamin A and E concentrations are also associated with dyslipidemia in these patients.

Key words: fat-soluble vitamins, oxidized LDL, prepubertal period

Streszczenie

Wprowadzenie: Produkty peroksydacji lipidów powstają w czasie fizjologicznych przemian zachodzących w komórkach. Jednakże, znacznie częściej pojawiają się w stanach patologicznych, gdy nadmiar wolnych rodników reaguje z nienasyconymi kwasami tłuszczowymi szczególnie lipoprotein o niskiej gęstości (LDL).

Celem pracy było zbadanie zależności pomiędzy oksydowanymi LDL (oxLDL), staniem witamin rozpuszczalnych w tłuszczach a parametrami antropometrycznymi u dzieci otyłych w okresie przedpokwitaniowym.

Materiał i metody: Grupę badaną stanowiło 32 dzieci otyłych (SDS-BMI >2) oraz 20 dzieci z prawidłową masą ciała (SDS-BMI <-1+1>). Stężenie oxLDL oznaczano w surowicy za pomocą testu ELISA. Witaminę A i E mierzono metodą wysokociśnieniowej chromatografii cieczowej. Poziom całkowitego cholesterolu, cholesterolu LDL i HDL oraz poziom triglicerydów oznaczano metodami enzymatycznymi.

Wyniki: Stężenie oxLDL i witaminy A było wyższe u dzieci otyłych niż w grupie kontrolnej odpowiednio o około 50% ($p=0,01$) i 40% ($p=0,001$). U dzieci otyłych stwierdzono istotną dodatnią korelację między stężeniem oxLDL a stężeniem witaminy A ($p<0,05$). Ponadto, stężenie oxLDL korelowało pozytywnie z wartościami BMI ($p<0,05$) i ilością tkanki tłuszczowej (kg) ($p<0,02$) u tych pacjentów. Stężenia witamin A i E korelowały z poziomem cholesterolu całkowitego (odpowiednio $p<0,05$, $p<0,01$). Oprócz tego zaobserwowano dodatnią korelację pomiędzy witaminą E i cholesterolem frakcji LDL ($p<0,05$).

Wnioski: Wstępne badanie wykazały, że wzrost stężenia oxLDL pojawia się u dzieci otyłych już w okresie przedpokwitaniowym i może poprzedzać u nich zmiany miażdżycowe. Uzyskane wyniki sugerują występowanie zależności pomiędzy witaminą A i oxLDL u tych pacjentów. Stwierdzono też związki pomiędzy stężeniem witaminy A i E we krwi a dyslipidemią u dzieci otyłych w okresie przedpokwitaniowym.

Słowa kluczowe: witaminy rozpuszczalne w tłuszczach, oksydowane LDL, okres przedpokwitaniowy

DEV PERIOD MED. 2017;XXI,3:266-271

INTRODUCTION

According to many studies, oxidant-antioxidant balance is needed to prevent freeradical-induced damage in nucleic acids, proteins and lipid structures. An important consequence of disturbances in the oxidant-antioxidant balance is lipid peroxidation [1]. Although lipid peroxidation products are formed during normal cell metabolism, these appear mostly in pathological conditions via producing an excess of free radicals that can react with unsaturated fatty acids, in particular low-density lipoprotein (LDL) [2]. In adults, higher oxLDL concentrations have been associated with higher body mass index (BMI), body fat, and insulin resistance [3-5]. Oxidation of this protein is considered a key factor in the pathogenesis of the metabolic syndrome, atherosclerosis and cardiovascular disease (). Elevated oxidized LDL (oxLDL) concentrations were observed mostly in obese adults, but much less is known about its relation to obesity in childhood [6-8]. In addition, obesity in adults and children is associated with an inflammatory condition that correlates with oxidative stress and may cause greater utilization of antioxidants [9].

Fat-soluble vitamins, which act as free radical scavengers, belong to the most important components of the antioxidant defense of lipids. It is suggested that the process of reducing oxidative damages may require higher concentrations of antioxidants, such as vitamin A and vitamin E [10]. Vitamin A (retinol) is a micronutrient required for growth and development, conceivably affecting lipid metabolism, energy regulation and body composition [11]. Vitamin E is most commonly found in the form of alpha-tocopherol and acts as a peroxy scavenger and thus prevents the oxidation of membrane lipids [12]. Given

the importance of these vitamins in lipid metabolism, there is also a need to investigate the relationship between fat-soluble vitamins and lipid peroxidation products in obesity. Recent studies in young populations suggest that fat deposition and chronic inflammation are associated with deficiencies in concentrations of fat-soluble vitamins [13-15]. Other studies show positive relations between levels of these vitamins and measures of obesity [16]. Thus, research data concerning these associations in obese children are inconsistent.

Therefore, the aim of this study was to investigate the relations between oxidized low-density lipoproteins, fat-soluble vitamin status and anthropometric parameters in prepubertal obese children.

METHODS

Thirty-two obese children (age, 7.8 ± 1.5 years; boys, 47%) were included in the study. Healthy normal-weight children ($n=25$) were the reference group. Obese and non-obese children who were taking medications that could affect growth, pubertal development, nutritional status or dietary intake were not included. Physical examination was performed and body mass index (BMI) was calculated. Children were classified as obese (SDS-BMI >2) and non-obese (SDS-BMI <-1+1>). Body composition was measured by dual-energy X-ray absorptiometry (DXA) using Lunar Prodigy (General Electric Healthcare, UK) with pediatric software 9.30.044.

We assessed the average daily energy intake and the percentage of energy intake from protein, fat and carbohydrates in the diets of obese and non-obese children. Patients had 3-5 meals every day. The dietary intake and physical activity data were collected using randomly

selected 3-day records. Average daily food rations and their nutritional value were calculated using nutritional analysis software (Dieta5) [17].

Venous blood samples were collected between 8.00 and 10.00 in the morning after overnight fasting and centrifuged (1000g for 10 min at 4°C). The concentration of oxLDL was determined in the serum by the ELISA assay (Immundiagnostik AG, Germany). Vitamins A and E in the serum were determined by the high-pressure liquid chromatography method (HPLC), (Knauer, Germany) [18]. Total cholesterol, LDL- and HDL-cholesterol and triglyceride levels were measured by enzymatic methods (ROCHE, Switzerland).

This study had been approved by the Ethics Committee of the Institute of Mother and Child. Informed consent was obtained from the parents of all the examined children.

Statistical analysis was performed using Statistica version 8.0. The results are presented as mean±standard deviation (SD) for normally distributed data or medians and interquartile range (25th-75th percentiles) for non-normally distributed variables. The Kolmogorov-Smirnov test and graphical inspections of data were used for evaluating the distribution for normality. Differences in anthropometric characteristics and biochemical parameters of obese and non-obese children were assessed using the Student's t-test for normally distributed data and the non-parametric Mann-Whitney test for non-normally distributed variables. Spearman or Pearson correlations between parameters were calculated. Data are presented with $p < 0.05$ considered statistically significant.

RESULTS

As expected, there are significant differences in the body weight, BMI and body composition between obese and non-obese children of the same age (tab. I). Average BMI was about 50% ($p < 0.001$) higher in obese patients than in controls. The obese children had a fourfold greater ($p < 0.001$) fat mass and 2-fold higher percentage of fat mass ($p < 0.001$) compared with non-obese subjects. Moreover, obese patients had greater lean tissue mass and total BMC by approximately 30% ($p < 0.001$). The daily energy intake of obese children before intervention was higher ($p < 0.05$) compared with the controls. However, the proportions of protein, fat and carbohydrates in the daily energy intake were similar in both groups (tab. I).

The concentrations of oxLDL and vitamin A were higher in obese children than in normal-weight controls by about 50% ($p = 0.01$) and 40% ($p = 0.001$), respectively (tab. II). No difference was observed in the concentration of vitamin E between both groups. The level of HDL-cholesterol was lower ($p = 0.001$), while the level of triglyceride was higher ($p = 0.01$) in obese children than in controls.

A significant positive correlation was found between oxLDL and vitamin A concentrations ($p < 0.05$) in obese children (tab. III). In addition, oxLDL correlated positively with BMI values ($p < 0.05$) and the amount of fat mass (kg) ($p < 0.02$) in these patients. No correlations were observed between oxLDL and lipid parameters.

Concentrations of vitamins A and E correlated with the level of total cholesterol ($p < 0.05$; $p < 0.01$, respectively). In addition, vitamin E correlated positively with LDL-cholesterol ($p < 0.05$). No correlations were observed between fat-soluble vitamins and triglycerides, or HDL-cholesterol.

DISCUSSION

Oxidized low-density lipoprotein is one that may reflect the oxidative stress in organisms [19]. An increase in oxLDL concentrations was found in populations of obese children with a wide age range and different pubertal stages [7-9, 20]. In this study, we showed higher concentrations of oxLDL associated with dyslipidemia including lower concentrations of HDL-cholesterol and higher concentrations of triglyceride in prepubertal obese children compared with normal-weight subjects. Similarly to our results, Okur et al. [8] observed higher values of oxidized protein related with disturbances in the lipid profile in obese children during the same pubertal stage. In addition, we observed positive relations between oxLDL and BMI, as well as the amount of fat mass (kg), but not the percentage of fat mass (%). It is suggested that greater adiposity, especially abdominal fat and low-extremity adiposity is associated with increased levels of oxidative stress. Furthermore, higher waist circumference, leptin and insulin resistance have been significantly linked with higher oxLDL levels in pediatric populations [8, 21]. These changes might be partially reversible by lifestyle intervention, even if patients do not reach normal body weight [22].

According to many authors, greater adiposity is associated with increased levels of oxidative stress, leading to reduced concentrations of antioxidants, including fat-soluble vitamins [9, 23]. These associations may result from differences in the energy intake and/or sequestering of lipophilic vitamins in adipose tissue [24]. Lower values of fat-soluble vitamins in American and European populations of obese children compared with normal-weight controls were observed [25, 26]. In addition, it is suggested that the increased production of reactive oxygen species in obesity may lead to increased utilization and thus low availability of antioxidant enzymes and vitamins [27].

In opposition to these studies, we found higher concentrations of vitamin A in obese than in normal-weight children, but these values are still within the reference range (0.8-2.8 $\mu\text{mol/l}$) [28]. A significant association between serum retinol concentrations and adiposity was found by Gunanti et al. [15] in Mexican-American children 8-15 years of age. Garcia et al. [16] also observed positively associated vitamin A with BMI and abdominal fat mass in Mexican school-age children. In addition, serum retinol concentrations were elevated in overweight and obese Brazilian adolescents, while no differences in vitamin E concentrations were found between these groups [29]. It seems that the homeostatic regulation of vitamin A through storage in the liver and controlled release from the liver may provide tissue with optimal amounts of retinol, in this way protecting

Table I. Anthropometric characteristics and dietary intake of obese and non-obese children.

Tabela I. Charakterystyka wybranych cech antropometrycznych i udział podstawowych składników pokarmowych w średniej racji pokarmowej u dzieci otyłych i w grupie kontrolnej.

	Obese children <i>Dzieci otyłe</i>	Non-obese children <i>Dzieci z prawidłową masą ciała</i>	P-value <i>P</i>
Age (y) ^a <i>Wiek (lata)</i>	7.8 ± 1.2	7.5 ± 1.5	0.479
Girls/Boys <i>Dziewczynki/Chłopcy</i>	17/15	13/12	
Weight (kg) ^b <i>Masa ciała (kg)^b</i>	40.2 (36.7-49.7)	23.4 (19.8-26.7)	<0.001
BMI (kg/m ²) ^b <i>BMI (kg/m²)^b</i>	23.4 (21.1-24.7)	15.3 (14.5 -15.9)	<0.001
Body composition: <i>Skład ciała:</i>			
Fat mass (%) ^a <i>Masa tłuszczowa (%)^a</i>	40.7 ± 4.5	20.2 ± 6.3	<0.001
Fat mass (kg) ^b <i>Masa tłuszczowa (kg)^b</i>	15.8 (12.8-19.6)	4.0 (2.9-5.8)	<0.001
Lean mass (kg) ^a <i>Masa mięśniowa (kg)^a</i>	24.5 ± 3.0	18.4 ± 3.0	<0.001
Total BMC (kg) ^b <i>Całkowite BMC (kg)^b</i>	1.2 (1.0-1.4)	0.9 (0.7 -1.0)	<0.001
Dietary intake: <i>Udział podstawowych</i> <i>składników pokarmowych:</i>			
Energy (kcal/d) ^b <i>Energia (kcal/d)^b</i>	1745 (1331-2075)	1540 (1240-1824)	0.032
Protein (% of energy intake) ^b <i>Udział energii z białka (%)</i>	13.2 (12.1-15.0)	13.8 (11.5-14.6)	0.445
Carbohydrate (% of energy intake) ^b <i>Udział energii z węglowodanów (%)</i>	53.5 (49.0-56.4)	52.2 (47.2-56.3)	0.548
Fat (% of energy intake) ^b <i>Udział energii z tłuszczu (%)</i>	33.3 (30.6-36.8)	34.0 (29.4-39.5)	0.939

^aData are presented as mean values±SD; *Dane przedstawione jako wartości średnie±SD*^bData are presented as median values (25th-75th percentiles); *Dane przedstawione jako mediany (25-75 percentyl)*
BMC, bone mineral content (*zawartość minerałów w kości*)

Table II. Biochemical measurements in obese and non-obese children.

Tabela II. Parametry biochemiczne u dzieci otyłych i w grupie kontrolnej.

	Obese children <i>Dzieci otyłe</i>	Non-obese children <i>Dzieci z prawidłową masą ciała</i>	P-value <i>P</i>
oxLDL(ng/ml) ^b	466 (313 – 658)	303(167-438)	0.010
Vitamin A (µmol/l) ^a	1.62 ± 0.46	1.15 ± 0.13	0.001
Vitamin E (µmol/l) ^a	18.5 ± 5.24	17.1 ± 1.76	0.238
LDL-cholesterol (mg/dl) ^b	104(94-117)	105(95-118)	0.799
HDL-cholesterol(mg/dl) ^a	49.7 ± 8.4	62.0 ± 9.3	0.001
Total-cholesterol(mg/dl) ^a	171 ± 20	177 ± 20	0.368
Triglycerides (mg/dl) ^b	79.0(59.8-129)	56.5(42.8-77.8)	0.010

^aData are presented as mean values±SD; *Dane przedstawione jako wartości średnie±SD*^bData are presented as median values (25th-75th percentiles); *Dane przedstawione jako mediany (25-75 percentyl)*

Table III. Relationships between biochemical and anthropometric parameters in obese children.

Tabela III. Zależności pomiędzy parametrami biochemicznymi i antropometrycznymi u dzieci otyłych.

	BMI BMI	Fat mass (%) Masa tłuszczowa	Fat mass (kg) Masa tłuszczowa	Total-cholesterol Całkowity cholesterol	LDL-cholesterol LDL-cholesterol	oxLDL oxLDL
oxLDL	0.360*	0.330	0.408**	-0.055	-0.008	-
Vitamin A	-0.003	0.040	0.030	0.381*	0.331	0.348*
Vitamin E	-0.280	-0.220	-0.257	0.448***	0.386*	0.128

*p<0.05; **p<0.02; ***p<0.01

against the toxicity of this vitamin [15]. However, it is not impossible that the process of reduction of oxidative damages requires increased antioxidant concentrations, among them fat-soluble vitamins.

We observed positive correlations between oxLDL and vitamin A in prepubertal obese children. However, no relations between oxidized protein and vitamin E were obtained in these patients. Similar to us, Beck et al. [30] observed higher values of oxLDL in obese women, and did not find any association between oxidized LDL and vitamin E in obese subjects. Moreover, Albuquerque et al. [29] found that elevated serum retinol but not alpha-tocopherol concentrations were associated with dyslipidemia in overweight or obese adolescents. Relations between both vitamins and lipid parameters were showed by Garcia et al. [16]. These authors found positive relations in obese children between fat-soluble vitamins and triglycerides, as well as total cholesterol. In our previous and in presented studies we found disturbances in body composition and abnormal lipid profile in prepubertal obese children (31, 32). In this study we also found positive relations between vitamins A and E and total cholesterol and additionally between vitamin E and LDL-cholesterol in obese patients. We agree that higher concentrations of lipids may require higher concentrations of fat-soluble vitamins to prevent deficiencies of these vitamins and increased risk of cardiovascular diseases [16].

CONCLUSION

Our preliminary study shows that oxLDL starts early during the prepubertal period and may precede atherosclerotic lesions. We suggest the occurrence of relationships between vitamin A and oxidized LDL in prepubertal obese children. Vitamin A and E concentrations are also associated with dyslipidemia in these patients.

REFERENCES

- Chiavaroli V, Giannini C, De Marco S, Chiarelli F, Mohn A. Unbalanced oxidant-antioxidant status and its effects in pediatric diseases. *Redox Report* 2011;16:101-107.
- Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *The New England Journal of Medicine* 2005;352:1685-1695.
- Neuparth MJ, Proença JB, Santos-Silva A, Coimbra S. Adipokines, oxidized low-density lipoprotein, and C-reactive protein levels in lean, overweight, and obese Portuguese patients with type 2 diabetes. *ISRN Obesity*, 2013, doi: 10.1155/2013/142097.
- Babakr AT, Elsheikh OM, Almarzouki AA, Assiri AM, Abdalla BE, Zaki HY, Fatani SH, NourEldin EM. Relationship between oxidized low-density lipoprotein antibodies and obesity in different glycemic situations. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy* 2014;7:513-520.
- Park JH, Park H, Lim ST, Park JK. Effects of a 12-week healthy-life exercise program on oxidized low-density lipoprotein cholesterol and carotid intima-media thickness in obese elderly women. *Journal of Physical Therapy Science* 2015;27(5):1435-1439.
- Kelly AS, Jacobs DR Jr, Sinaiko AR, Moran A, Steffen LM, Steinberger J. Relation of circulating oxidized LDL to obesity and insulin resistance in children. *Pediatric Diabetes* 2010;11(8):552-555.
- Norris AL, Steinberger J, Steffen LM, Metzger AM, Schwarzenberg SJ, Kelly AS. Circulating oxidized LDL and inflammation in extreme pediatric obesity. *Obesity (Silver Spring)* 2011;19(7):1415-1419.
- Okur I, Tumer L, Ezgu FS, Yesilkaya E, Aral A, Oktar SO, Bideci A, Hasanoglu A. Oxidized low-density lipoprotein levels and carotid intima-media thickness as markers of early atherosclerosis in prepubertal obese children. *Journal of Pediatric Endocrinology and Metabolism* 2013;26(7-8):657-662.
- Vincent HK, Innes KE, Vincent KR. Oxidative stress and potential interventions to reduce oxidative stress in overweight and obesity. *Diabetes, Obesity and Metabolism* 2007;9:813-839.
- Mikoluc B, Motkowski R, Karpinska J, Piotrowska-Jastrzebska J. Plasma levels of vitamins A and E, coenzyme Q10, and anti-ox-LDL antibody titer in children treated with an elimination diet due to food hypersensitivity. *International Journal for Vitamin and Nutrition Research* 2009;79:328-336.
- Kiefer FW, Orasanu G, Nallamshetty S, Brown JD, Wang H, Luger P, Qi NR, Burant CF, Duester G, Plutzky J. Retinaldehyde dehydrogenase 1 coordinates hepatic gluconeogenesis and lipid metabolism. *Endocrinology*. 2012;153:3089-3099.
- Niki E. Role of vitamin E as a lipid-soluble peroxyl radical scavenger: in vitro and in vivo evidence. *Free Radical Biology and Medicine* 2014;66:3-12.
- Sarni RO, Suano de Souza FI, Ramalho RA, Schoeps Dde O, Kochi C, Catherino P, Dias MC, Pessotti CF, Mattoso LC,

- Colugnat FA. Serum retinol and total carotene concentrations in obese pre-school children. *Medical Science Monitor* 2005;11:CR510-514.
14. Wei X, Peng R, Cao J, Kang Y, Qu P, Liu Y, Xiao X, Li T. Serum vitamin A status is associated with obesity and the metabolic syndrome among school-age children in Chongqing, China. *Asia Pacific Journal of Clinical Nutrition* 2016;25:563-570.
 15. Gunanti IR, Marks GC, Al-Mamun A, Long KZ. Low serum concentrations of carotenoids and vitamin E are associated with high adiposity in Mexican-American children. *The Journal of Nutrition* 2014;144:489-495.
 16. García OP, Ronquillo D, del Carmen Caamaño M, Martínez G, Camacho M, López V, Rosado JL. Zinc, iron and vitamins A, C and E are associated with obesity, inflammation, lipid profile and insulin resistance in Mexican school-aged children. *Nutrients* 2013;5:5012-5030.
 17. Wajszczyk B, Chwojnowska Z, Nasiadko D, Rybaczuk M. Dieta 5.0 software for individual and group nutrition assessment and diet planning. National Food and Nutrition Institute, Warsaw, Poland, 2015.
 18. Zaman Z, Fielden P, Frost PG. Simultaneous determination of vitamins A and E and carotenoids in plasma by reversed-phase HPLC in elderly and younger subjects. *Clinical Chemistry* 1993;39:2229-2234.
 19. Itabe H, Ueda M. Measurement of plasma oxidized low-density lipoprotein and its clinical implications. *Journal of Atherosclerosis and Thrombosis* 2007;14:1-11.
 20. Matusik P, Prokopowicz Z, Norek B, Olszanecka-Glinianowicz M, Chudek J, Malecka-Tendera E. Oxidative/Antioxidative status in obese and sport trained children: a comparative study. *Biomed Research International* 2015; doi: 10.1155/2015/315747.
 21. Kelishadi R, Hashemi M, Mohammadifard N, Asgary S, Khavarian N. Association of changes in oxidative and proinflammatory states with changes in vascular function after a lifestyle modification trial among obese children. *Clinical Chemistry* 2008;54:147-153.
 22. Morell-Azanza L, García-Calzón S, Rendo-Urteaga T, Martín-Calvo N, Chueca M, Martínez JA, Azcona-Sanjulián MC, Martí A. Serum oxidized low-density lipoprotein levels are related to cardiometabolic risk and decreased after a weight loss treatment in obese children and adolescents. *Pediatric Diabetes* 2016; doi: 10.1111/pedi.12405.
 23. Strauss RS. Comparison of serum concentrations of alpha-tocopherol and beta-carotene in a cross-sectional sample of obese and nonobese children (NHANES III). *National Health and Nutrition Examination Survey. The Journal of Pediatrics* 1999;134:160-165.
 24. Landrier JF, Marcotorchino J, Tourniaire F. Lipophilic micronutrients and adipose tissue biology. *Nutrients* 2012;4:1622-1649.
 25. Ford ES, Gillespie C, Ballew C, Sowell A, Mannino DM. Serum carotenoid concentrations in US children and adolescents. *The American Journal of Clinical Nutrition* 2002;76:818-827.
 26. Mohn A, Catino M, Capanna R, Giannini C, Marcovecchio M, Chiarelli F. Increased oxidative stress in prepubertal severely obese children: effect of a dietary restriction-weight loss program. *Journal of Clinical Endocrinology and Metabolism* 2005;90:2653-2658.
 27. Murer SB, Aeberli I, Braegger CP, Gittermann M, Hersberger M, Leonard SW, Taylor AW, Traber MG, Zimmermann MB. Antioxidant supplements reduced oxidative stress and stabilized liver function tests but did not reduce inflammation in a randomized controlled trial in obese children and adolescents. *The Journal of Nutrition* 2014;144:193-201.
 28. Prószyńska K, Lorenc RS. Witaminy A,D,E znaczenie, zapotrzebowanie, suplementacja, toksyczność. *Medycyna* 2000 1992;3: 40-43.
 29. Albuquerque MN, Diniz Ada S, Arruda IK. Elevated serum retinol and low beta-carotene but not alpha-tocopherol concentrations are associated with dyslipidemia in Brazilian adolescents. *Journal of Nutritional Science and Vitaminology (Tokyo)* 2016;62:73-80.
 30. Beck J, Ferrucci L, Sun K, Fried LP, Varadhan R, Walston J, Guralnik JM, Semba RD. Circulating oxidized low-density lipoproteins are associated with overweight, obesity, and low serum carotenoids in older community-dwelling women. *Nutrition* 2008;24:964-968.
 31. Gajewska J, Klemarczyk W, Ambroszkiewicz J, Chełchowska M, Riahi A, Zielińska A, Oltarzewski M, Laskowska-Klita T. Effect of weight reduction programme on C-peptide concentration and lipid profile in obese children aged 4 to 10 years. *Medycyna Wieku Rozwojowego* 2010;14:357-364.
 32. Gajewska J, Weker H, Ambroszkiewicz J, Chełchowska M, Więch M, Laskowska-Klita T. Changes in concentration of serum adiponectin multimeric forms following weight reduction programme in prepubertal obese children. *Medycyna Wieku Rozwojowego* 2011;15:298-305.

Author's contributions/Wkład Autorów

According to the order of the Authorship/Według kolejności

Conflicts of interest/Konflikt interesu

The Authors declare no conflict of interest.
Autorzy pracy nie zgłaszają konfliktu interesów.

Received/Nadesłano: 07.06.2017 r.

Accepted/Zaakceptowano: 20.06.2017 r.

Published online/Dostępne online

Address for correspondence:

Joanna Gajewska

Screening Department

Institute of Mother and Child

Kasprzaka 17a, 01-211 Warsaw

tel, (048 22) 32-77-260,

e-mail: joanna.gajewska@imid.med.pl